



FIGURE 1



**FIGURE 2**


Anti-IgE Mab binding to IgE or receptor bound IgE



bioRxiv preprint doi: <https://doi.org/10.1101/092466>; this version posted May 1, 2016. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.


### FIGURE 3

Ability of Recombinant Exon 3 to Inhibit Conjugated 8H8 (5 ug/ml) or 15A.2 (2.5 ug/ml) Mab from binding to an IgE solid phase




## FIGURE 4

Ability of soluble, purified Canine IgE to inhibit Mab 8H8 ( 2.5 ug/ml) or 15A.2 ( 2.5 ug/ml ) from binding to an IgE solid phase



00000-094800

FIGURE 5



**FIGURE 6**


<b>LIBRARY</b>	<b>SEQUENCE OF DISPLAYED PEPTIDE</b>	<b>ISOLATE</b>
PhDc7c	C S N P H V T H C	M13 1&9
PhDc7c	C S H P H L T H C	M13 7
PhDc7c	C S N P H I T Q C	M13 10
PhDc7c	C M N P H I T H C	M13 14
PhDc7c	C T N P H N P Y C	M13 2&8
PhDc7c	C P N P H N P Y C	M13 3&5
PhD12	V T L C P N P H I P M C	M13 48
PhDc7c	C H P H L P K S C	M13 4&12
PhDc7c	C H P H L P K R C	M13 6
	Y C R V T H P H L P K D I V R S I	Canine IgE

## FIGURE 7

## SPECIES

Canine  
Human  
Green Monkey  
Cat  
Swine  
Mouse  
Horse

FIGURE 8





**FIGURE 9**




FIGURE 10A





FIGURE 10B



000000 09470000

FIGURE 11

Mab conjugate titration on purified canine IgE solid phase




FIGURE 12




FIGURE 13




FIGURE 14





FIGURE 15



000000-09270000

FIGURE 16




FIGURE 17

